

WO9704794

Publication Title:

MODULATORS OF EXPRESSION AND FUNCTION OF LRP IN ALZHEIMER'S DISEASE

Abstract:

Abstract of WO9704794

The present invention broadly relates to the treatment, diagnosis, and prophylactic prevention of Alzheimer's disease. More specifically, the present invention relates to methods and compositions for preventing the endocytosis and cellular internalization of integral membrane amyloid beta -precursor protein (APP) and its subsequent catabolism by blocking or interfering with the association or binding of APP with members of the low density lipoprotein receptor family. Data supplied from the esp@cenet database - Worldwide

Courtesy of <http://v3.espacenet.com>

This Patent PDF Generated by Patent Fetcher(TM), a service of Stroke of Color, Inc.

ATTORNEY DOCKET NUMBER: 8449-304-999
SERIAL NUMBER: 10/530,393
REFERENCE: B04

BEST AVAILABLE COPY

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 37/18, 38/00, 39/395, C07K 1/00, 14/00, 16/00, 17/00, C12P 21/04, G01N 33/53, 33/567	A1	(11) International Publication Number: WO 97/04794 (43) International Publication Date: 13 February 1997 (13.02.97)
(21) International Application Number: PCT/US96/12686 (22) International Filing Date: 29 July 1996 (29.07.96) (30) Priority Data: 60/001,600 27 July 1995 (27.07.95) US 60/001,653 28 July 1995 (28.07.95) US (71) Applicants: THE AMERICAN NATIONAL RED CROSS [US/US]; 430 17th Street, N.W., Washington, DC 20006 (US). THE GENERAL HOSPITAL CORPORATION [US/US]; Fruit Street, Boston, MA 02114 (US). (72) Inventors: STRICKLAND, Dudley, K.; Olney, MD (US). HYMAN, Bradley, T.; Swampscott, MA (US). KOUNNAS, Maria, Z.; Gaithersburg, MD (US). MOIR, Robert, D.; Boston, MA (US). TANZI, Rudolph, E.; Canton, MA (US). (74) Agents: GOLDSTEIN, Jorge, A. et al.; Sterne, Kessler, Goldstein & Fox P.L.L.C., Suite 600, 1100 New York Avenue, N.W., Washington, DC 20005-3934 (US).		(81) Designated States: CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: MODULATORS OF EXPRESSION AND FUNCTION OF LRP IN ALZHEIMER'S DISEASE		
(57) Abstract The present invention broadly relates to the treatment, diagnosis, and prophylactic prevention of Alzheimer's disease. More specifically, the present invention relates to methods and compositions for preventing the endocytosis and cellular internalization of integral membrane amyloid β -precursor protein (APP) and its subsequent catabolism by blocking or interfering with the association or binding of APP with members of the low density lipoprotein receptor family.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Modulators of Expression and Function of LRP in Alzheimer's Disease

Rights of the United States Government

Part of the work performed during development of this invention utilized U.S. government funds in the form of grant GM42581, grant HL50787, and grant AG12406, all from the National Institutes of Health. The U.S. government therefore has certain rights in one or more embodiments of this invention.

Field of the Invention

The present invention broadly relates to the treatment, diagnosis, and prophylactic prevention of Alzheimer's disease. More specifically, the present invention relates to methods and compositions for preventing the endocytosis and cellular internalization of integral membrane amyloid β -precursor protein (APP) and its subsequent catabolism by blocking or interfering with the association or binding of APP with members of the low density lipoprotein receptor family.

Background of the Invention

Alzheimer's disease (AD) is a late onset neurodegenerative disorder characterized by the extracellular deposition of insoluble aggregates composed of the 40 to 42 amino acid A β peptide in the brain (Glennner and Wong, *Biochem. Biophys. Res. Commun.* 120:885-890 (1984); Masters *et al.*, *EMBO J.* 4:2757-2763 (1985)). A β peptide is derived from an integral membrane protein termed amyloid β -protein precursor protein (APP) (Tanzi *et al.*, *Science* 235:880 (1987); Kang *et al.*, *Nature* 325:733-736 (1987)). The function and metabolism of APP have been the subject of intensive study due to the fact that mutations in APP are associated with an autosomal dominant form of AD, (Goate *et al.*, *Nature* 349:704-707 (1991)) and over-production of APP is the presumptive cause of AD in trisomy 21 (Tanzi *et al.*, *Science* 235:880 (1987); Hyman *et al.*, *Proc. Natl. Acad. Sci. USA* 92:3586-3590 (1995)). Multiple APP isoforms can be generated by alternatively splicing of mRNAs. The major isoforms in brain are APP695, APP751, and APP770 containing 695, 751 and 770 amino acids, respectively. These isoforms are transmembranous proteins having large extracellular regions,

with hydrophobic membrane spanning domains and short cytoplasmic segments. APP is also a member of an evolutionary conserved family of proteins which include the APP-like proteins, APLP1 and APLP2 (Wasco *et al.*, *Proc. Natl. Acad. Sci. USA* 89:10758-10762 (1992); Wasco *et al.*, *Nature Genet.* 5:95-100 (1993); Slunt *et al.*, *J. Biol. Chem.* 269:2637-2644 (1994)).

Secreted forms of APP are generated by proteolytic cleavages within their extracellular domain close to the transmembrane region. The extracellular regions of APP751, APP770, and APLP2 each contain a Kunitz protease inhibitor (KPI) domain encoded by an alternatively-transcribed exon (Kitaguchi *et al.*, *Nature* 331:530-532 (1988); Tanzi *et al.*, *Nature* 331:528-530 (1988); Wasco *et al.*, *Nature Genet.* 5:95-100 (1993); Slunt *et al.*, *J. Biol. Chem.* 269:2637-2644 (1994)). Secreted forms of APP having the KPI domain correspond to a protease inhibitor that has been identified separately and named protease nexin II (APP/PN-2) (Van Nostrand and Cunningham, *J. Biol. Chem.* 262:8508-8514 (1987); Oltersdorf *et al.*, *Nature* 341:144-147 (1989); Van Nostrand *et al.*, *Nature* 341:546-549 (1989)), a potent inhibitor of the blood coagulation factors IXa (Schmaier *et al.*, *J. Clin. Invest.* 92:2540-2545 (1993)) and XIa (Van Nostrand *et al.*, *J. Biol. Chem.* 265:9591-9594 (1990)). APP/PN-2 binds with high affinity to cultured fibroblasts (Johnson-Wood *et al.*, *Biochem. Biophys. Res. Commun.* 200:1685-1692 (1994)), and APP/PN-2:proteinase complexes are internalized and degraded by cultured cells (Knauer and Cunningham, *Proc. Natl. Acad. Sci. USA* 79:2310-2314 (1982); Knauer *et al.*, *J. Cell. Physiol.* 117:385-396 (1983)) although the mechanism for this process is unknown. Recent studies have identified the low density lipoprotein receptor-related protein (LRP) as the receptor responsible for the catabolism of another Kunitz-type inhibitor, tissue factor pathway inhibitor (TFPI) (Warshawsky *et al.*, *Proc. Natl. Acad. Sci. USA* 91:6664-6668 (1994)).

LRP is a large multiligand receptor (Krieger and Herz, *Annu. Rev. Biochem.* 63:601-637 (1994)) that is a member of the LDL receptor family, which also includes the LDL receptor (Yamamoto *et al.*, *Cell* 39:27-38 (1984)), the VLDL receptor (Takahashi *et al.*, *Proc. Natl. Acad. Sci. USA* 89:9252-9256

(1992)), and glycoprotein 330 (Saito *et al.*, *Proc. Natl. Acad. Sci. USA* 91:9725-9729 (1994)). A 39 kDa protein, termed the receptor associated protein (RAP) (Strickland *et al.*, *J. Biol. Chem.* 266:13364-13369 (1991)) binds to members of the LDL receptor family (Williams *et al.*, *J. Biol. Chem.* 267:9035-9040 (1992); Kounnas *et al.*, *J. Biol. Chem.* 267:21162-21166 (1992); Battey *et al.*, *J. Biol. Chem.* 269:23268-23273 (1994)) and blocks their ligand binding capacity. LRP mediates the cellular uptake and subsequent degradation of proteinases, such as tissue-type plasminogen activator (Bu *et al.*, *Proc. Natl. Acad. Sci. USA* 89:7427-7431 (1992)) and urokinase-type plasminogen activator (Kounnas *et al.*, *J. Biol. Chem.* 268:21862-21867 (1993)), proteinase-inhibitor complexes, such as α_2 -macroglobulin-proteinase complexes (Ashcom *et al.*, *J. Cell Biol.* 110:1041-1048 (1990); Moestrup and Gliemann, *J. Biol. Chem.* 264:15574-15577 (1989)), serpin-proteinase complexes (Orth *et al.*, *Proc. Natl. Acad. Sci. USA* 89:7422-7426 (1992); Nykjaer *et al.*, *J. Bio. Chem.* 267:14543-14546 (1992); Poller *et al.*, *J. Biol. Chem.* 270:2841-2845 (1995)), matrix proteins, such as thrombospondin (Mikhailenko *et al.*, *J. Biol. Chem.* 270:9543-9549 (1995)), apolipoprotein E (apoE)-enriched lipoproteins (Kowal *et al.*, *J. Biol. Chem.* 265:10771-10779 (1990); Beisiegel *et al.*, *Nature* 341:162-164 (1989)), hepatic lipase (Kounnas *et al.*, *J. Biol. Chem.* 270:9307-9312 (1995)) and lipoprotein lipase (Chappell *et al.*, *J. Biol. Chem.* 268:14168-14175 (1993)).

LRP is expressed in many tissues and is a major apoE receptor in the central nervous system (Rebeck *et al.*, *Neuron* 11:575-580 (1993)). Genetic data implicate inheritance of the $\epsilon 4$ allele of apoE as a risk factor in AD (Strittmatter *et al.*, *Proc. Natl. Acad. Sci. USA* 90:1977-1981 (1993); Rebeck *et al.*, *Neuron* 11:575-580 (1993); Poirier *et al.*, *Lancet* 342:697-699 (1993); Saunders *et al.*, *Neurology* 43:1467-1472 (1993)). A possible involvement of LRP in AD is suggested in part by the observation that LRP, as well as apoE and other LRP ligands, decorate senile plaques (Rebeck *et al.*, *Ann. Neurol.* 37:211-217 (1995)).

The ability of LRP to mediate the cellular catabolism of TFPI, a KPI-containing protein, led to the investigation of the role of LRP in the catabolism of APP₇₇₀. The present inventors have found that LRP is capable of binding and

mediating the internalization and degradation of APP,770 as well as its complexes with proteinases.

Because catabolism of APP has been shown to generate the A β peptide, which is believed to be the causative agent of Alzheimer's Disease, there is a need
5 for compositions and methods which reduce the interaction, cellular internalization and subsequent catabolism of APP.

Summary of the Invention

Accordingly, it is an object of the present invention to provide agents
10 which bind to APP or LDL-receptor family members and reduce the interaction, cellular internalization, and subsequent catabolism of APP. Other objects, features and advantages of the present invention will be set forth in the detailed description of preferred embodiments that follows, and in part will be apparent from the description or may be learned by practice of the invention. These objects
15 and advantages of the invention will be realized and attained by the compositions methods particularly pointed out in the written description and claims hereof.

A first embodiment of the present invention therefore relates to agents which bind to the APP-binding site on the LRP particle (Group I agents) and agents which bind to the LRP-binding site found on APP (Group II agents).

20 An additional embodiment of the present invention relates to DNA molecules which encode peptides and antibodies that are Group I agents and/or Group II agents and to host organisms that have been transformed with the DNA molecules of the present invention.

The present invention also relates to processes for preparing DNA
25 molecules which encode a functional derivative or a fragment of LRP, RAP or APP. These processes can yield DNA sequences which are inserted into a vector DNA containing expression control sequences in such a way that the expression control sequences regulate the expression of the inserted DNA.

A further embodiment of the present invention relates to methods for
30 preparing polypeptides that are functional derivatives of LRP which comprise

taking the polypeptide from the native receptor molecule by enzymatic, such as proteolytic, or chemical, such as reductive, treatment.

An additional embodiment of the invention relates to a process for preparing Group I agents which are a functional derivative of APP or RAP which
5 comprise expressing a recombinant DNA molecule according to the invention.

The present invention further relates to Group II agents which are antibodies, or an antibody fragment containing the antigen binding domain, that bind to the LRP binding site found on APP.

Another embodiment of the present invention relates to processes for
10 preparing Group II agents which are a functional derivative of LRP which comprise expressing a recombinant DNA molecule according to the invention.

The present invention additionally relates to hybrid cell lines that secrete monoclonal antibodies against the LDL-receptor protein which interfere with APP attachment to the LRP receptor.

15 An additional embodiment of the present invention includes the use of Group I agents and/or Group II agents for qualitatively and/or quantitatively determining or purifying the presence of LRP which is found in a sample.

A further embodiment of the present invention includes a test kit for determining whether a polypeptide is a Group I agent and/or a Group II agent, this
20 kit comprising a carrier means having in close confinement therein one or more container means at least one of which contains an antibody that binds to the LRP binding site found on APP.

Another embodiment of the present invention relates to processes for preparing antibodies that bind to the LRP binding site found on APP, in which a
25 host animal is immunized with one or more polypeptides of Group I and/or Group II, the B-lymphocytes of these host animals are fused with myeloma cells, and a hybrid cell line secreting the monoclonal antibody is subcloned and cultivated.

An additional embodiment of the present invention relates to the use of Group I agents and/or Group II agents, or the native receptor molecules of the
30 LDL-receptor family or pharmaceutically suitable salts thereof, for the therapeutic or prophylactic treatment of the human body.

The invention also relates to methods for reducing the rate of onset or the severity of Alzheimer's disease, comprising administering to an animal, such as a human, one or more Group I agents and/or one or more Group II agents in an amount effective to reduce the rate of APP attachment to its receptor.

5 A further embodiment of the present invention relates to pharmaceutical compositions for therapeutic treatment of Alzheimer's disease, comprising one or more of Group I agents and/or one or more Group II agents and/or the native receptor molecule of the LDL-receptor family and a pharmaceutically acceptable carrier.

10 An additional embodiment of the invention relates to the use of LRP for inhibiting the binding of natural ligands to a member of the LDL-receptor family of proteins.

 An alternative embodiment of the present invention relates to methods for identifying substances which inhibit the binding of a ligand (RAP) or APP to a protein derived from the LDL-receptor family, comprising the steps of:

- 15 a) incubating the receptor, or a soluble form of the receptor, with RAP or APP in the presence of a potential inhibitor substance; and
- b) determining the extent of binding of RAP or APP to the receptor or receptor fragment.

20 Another embodiment of the present invention relates to methods for detecting receptors of the LDL-receptor family, comprising the steps of:

- a) incubating a substance derived from a fragment of RAP or APP which contains a binding activity for the receptor with a sample; and
- 25 b) determining the extent of binding of the RAP or APP material to the sample.

 A further embodiment of the present invention relates to methods for supplying a therapeutically active substance into a carrying cell, characterized in

30 that

- a) a fragment of RAP or APP with a binding activity on the LDL-receptor is coupled with the therapeutic substance; and
- b) the said material is added to the corresponding cell material, bound to the receptor and in this way the therapeutically active substance is introduced into the cell.

It is to be understood that both the foregoing general description and the following detained description are exemplary and explanatory only and are intended to provide further explanation of the invention as claimed.

Brief Description of the Figures

Figure 1 shows the catabolism of APP by fibroblasts containing LRP (panel A) compared to the absence of catabolism of APP by fibroblasts deficient in LRP (panel B) and the absence of catabolism of APP lacking the KPI domain by either LRP-containing or LRP-deficient fibroblasts (panels C and D). Note the inhibition of catabolism of APP by the addition of RAP in fibroblasts containing LRP (panel A).

Figure 2 shows the internalization and degradation of LDL by both LRP-containing and LRP-deficient fibroblasts (panels A and B) and the internalization and degradation of α_2 -macroglobulin by LRP-containing fibroblasts and the lack of both internalization and degradation of α_2 -macroglobulin by LDL-deficient fibroblasts (panels C and D).

Figure 3 shows the internalization and degradation of APP by LRP-containing fibroblasts (panels A and B) and the substantially lower level of internalization and degradation of APP by LRP-deficient fibroblasts (panels C and D). Note the inhibition of internalization of APP by the addition of heparin (panels A, B, C and D).

Figure 4 shows the inhibition of APP degradation by LRP-binding ligands in LRP-containing fibroblasts (panel A) and the lesser effect of these ligands on APP degradation by LRP-deficient fibroblasts (panel B).

Figure 5 shows the direct ability of APP to bind LRP as measured by enzyme-linked immunosorbant assay (ELISA) using anti-APP (panel A) as

compared to the amount of LRP available for binding (panel B). Note the inhibition of APP binding by RAP, an LRP ligand (panel C).

Figure 6 shows the enhancement of degradation of APP in LRP-containing fibroblasts by coupling with coagulation factor IXa (panel A) as compared to the degradation of APP in LRP-deficient fibroblasts (panel B). Note the inability of this complexing to avoid inhibition of degradation by RAP (panel A).

Figure 7 shows a slight inhibition of APP degradation in LRP-containing fibroblasts by competition with apolipoprotein E3/ β very low density lipoprotein or apolipoprotein E4/ β very low density lipoprotein (panel A) as compared to the same competition in LRP-deficient fibroblasts.

Detailed Description of the Preferred Embodiments

As used herein, the LDL-receptor family is defined as those proteins which are recognized in the art as being formed from four structurally-related cell surface receptors and which mediate the endocytosis of lipoproteins and other plasma proteins (Brown *et al.*, *Curr. Opin. Lipidology* 2:65-72 (1991)). These receptors share the following common features: cysteine-rich repeats which are responsible for ligand binding; cysteine-rich repeats of the epidermal growth factor-type (EGF-type); tyrosine-tryptophan-threonine-aspartate repeats; a single region spanning the membrane; and at least one internalizing signal (Willnow *et al.*, *J. Biol. Chem.* 267:26172-21180 (1992)).

As used herein, an agent is said to reduce the amount or rate of binding if the amount or rate of binding is less in the presence of the agent than when the agent is absent. Under conditions when the amount or rate of reduction is nearly complete, there will be an actual inhibition or total blocking of binding.

As used herein, the agents of the present invention, *i.e.* the Group I agents and the Group II agents, may be any composition of matter provided that it has the ability to bind to the APP-binding site on LRP (Group I) and/or the ability to bind to the LRP-binding site on APP (Group II). Suitable agents exhibiting these properties include, but are not limited to, peptides, antibodies,

carbohydrates, nucleic acids, vitamins, pharmaceutical agents, and the like, including derivatives thereof.

The agents of the present invention may be identified and/or prepared according to any of the methods and techniques known to those skilled in the art.

5 These agents, particularly peptide agents and antibody agents, may occur or be produced as monomer, dimers, trimers, tetrameres or multimers. Such multimers can be prepared using enzymatic or chemical treatment of the native receptor molecules or be prepared using recombinant techniques.

10 Preferably, the agents of the present invention are selected and screened at random or rationally selected or designed using protein modeling techniques.

For random screening, candidate agents are selected at random and assayed for their ability to reduce the amount or rate of binding of the amyloid β -precursor protein (APP) to the LDL receptor-related protein (LRP). Any of the suitable methods and techniques known to those skilled in the art may be
15 employed to assay candidate agents.

For rational selection or design, the agent is selected based on the configuration of the LRP binding site found on APP or the APP binding site found on the LRP. Any of the suitable methods and techniques known to those skilled in the art may be employed for rational selection or design. For example, one skilled
20 in the art can readily adapt currently available procedures to generate antibodies, peptides, pharmaceutical agents and the like capable of binding to a specific peptide sequence of LRP or APP. Illustrative examples of such available procedures are described, for example, in Hurby *et al.*, "Application of Synthetic Peptides: Antisense Peptides," in *Synthetic Peptides, A User's Guide*, W.H. Freeman, NY, pp. 289-307 (1992); Kaspczak *et al.*, *Biochemistry* 28:9230 (1989);
25 and Harlow, *Antibodies*, Cold Spring Harbor Press, NY (1990).

The agents of the present invention can alternatively be identified using modification of methods known in the art. For example, suitable peptide agents may be identified using the filter binding assay described by Mischak *et al.*
30 (Mischak *et al.*, *J. Gen. Virol.* 69:2653-2656 (1988) and Mischak *et al.*, *Virology* 163:19-25 (1988)), wherein the peptide is applied to a suitable membrane, such as

nitrocellulose, and the membrane is saturated with a detergent mixture in order to block any non-specific binding. The treated membrane is then incubated with labeled rhinovirus, *e.g.* with HRV2 labeled with ^{35}S -methionine, in order to check the specific binding. After washing and drying of the membrane, specific binding
5 can be visualized by autoradiography.

As noted above, the Group I agents of the present invention include those agents which bind directly to the APP binding site found on the LRP.

Additionally, the Group I agents of the present invention bind to the LRP in interfering proximity with the APP binding site or bind to the LRP in such a
10 manner so as to conformationally alter the APP binding site. Suitable Group I agents can therefore be first identified by their ability to bind the LRP and then by their ability to reduce the amount or rate at which APP binds to LRP. Illustrative examples of Group I agents of the present invention include, but are not limited to: soluble fragments of APP containing the KPI domain; anti-LRP antibodies;
15 soluble fragments of receptor associated protein (RAP); α_2 -macroglobulin:proteinase complexes; pregnancy zone protein (PZP):proteinase complexes; tissue-type plasminogen activator; pro-urokinase-type plasminogen activator; tissue factor pathway inhibitor; apolipoprotein E-enriched lipoproteins; lipoprotein lipase; hepatic lipase; thrombospondin; and lactoferrin.

Preferred Group I agents are based on and derived from the amino acid
20 sequence of the receptor associated protein (RAP). An especially preferred type of Group I agent is isolated RAP or a fragment thereof, such as a soluble fragment of RAP which contains the LRP binding site. Such agents act as competitive inhibitors of APP binding to its receptor *in vitro* as well as *in vivo*.

The preferred fragments of RAP are soluble under physiological
25 conditions. The C-terminus of these polypeptides can be shortened as desired, provided that the binding capacity for the LRP particle remains intact. The preferred amino acid sequence of RAP corresponds to the human protein. Suitable RAP sequences can also be derived from the amino acid sequence of
30 RAP isolated from other mammals or amphibia.

RAP, or a fragment thereof, may be produced using any of the methods and techniques known to those skilled in the art. For example, RAP can be purified from a source which naturally expresses the protein, can be isolated from a recombinant host which has been altered to express RAP or fragment thereof, or
5 can be synthesized using protein synthesis techniques known in the art. The skilled artisan can readily adapt a variety of techniques in order to obtain Group I peptide agents which contain the LRP binding site found on RAP.

The isolation of native RAP proteins is known, as described, for example, in Ashcom *et al.*, *J. Cell. Biol.* 110:1041-1048 (1990) and Jensen *et al.*, *FEBS Lett.* 255:275-280 (1989). In order to generate fragments of RAP which contains
10 the LRP binding site, isolated native protein may be converted by enzymatic and/or chemical cleavage to generate fragments of the whole protein, for example by reacting cell lines which express an RAP with an enzyme such as papain or trypsin or a chemical such as cyanogen bromide. Proteolytically active enzymes
15 or chemicals are preferably selected in order to release the extracellular receptor region. Fragments which contain the LRP binding site, especially fragments which are soluble under physiological conditions, can then be isolated using known methods.

Alternatively, RAP or a fragment of RAP may be expressed in a recombinant bacteria, as described, for example, in Williams *et al.*, *J. Biol. Chem.* 267:9035-9040 (1992) and Wurshawsky *et al.*, *J. Biol. Chem.* 269:3325-3330
20 (1994).

The Group II agents of the present invention include compositions which bind to the LRP binding site found on APP. Additionally, the Group II agents of the present invention include compositions that bind to APP in interfering
25 proximity to the LRP binding site. Suitable Group II agents can therefore be first identified by their ability to bind to APP and then by their ability to reduce the amount or rate at which APP binds to LRP. Illustrative examples of the Group II agents of the present invention include, but are not limited to, antibodies which
30 bind to the LRP binding site found on APP and soluble fragments of LRP.

Preferred Group II agents include antibodies and antibody fragments which are capable of binding to a residue found on APP and consequently act as a competitive inhibitor for LRP binding. The most preferred antibodies or antibody fragments of the present invention bind to an LRP-specific epitope in APP.

5 Antibodies, or other Group II agents such as anti-sense peptides, which bind to epitopes within this sequence reduce the amount or rate of APP binding to LRP.

The antibodies of the present invention include polyclonal and monoclonal antibodies, as well as antibody fragments and derivatives that contain the relevant antigen binding domain of the antibodies. Such antibodies or antibody
10 fragments are preferably used in the diagnostic and therapeutic embodiments of the present invention.

Suitable monoclonal and polyclonal antibodies may be prepared by any of the methods and techniques well known in the art, such as described in, for example, A.M. Campbell, *Monoclonal Antibody Technology: Laboratory
15 Techniques in Biochemistry and Molecular Biology*, Elsevier Science Publishers, Amsterdam, The Netherlands (1984) and Harlow; *Antibodies*, Cold Spring Harbor Press, NY (1989). For example, an antibody capable of binding to a domain of APP can be generated by immunizing an animal with a polypeptide whose sequence is encoded by that domain. Any animal (mouse, rabbit, etc.) which is
20 known to produce antibodies can be utilized to produce antibodies with the desired specificity and suitable methods for immunization of these animals are well known in the art, including, for example, subcutaneous or intraperitoneal injection of the polypeptide. One skilled in the art will recognize that the amount of polypeptide used for immunization will vary based on a number of factors,
25 including the animal which is immunized, the antigenicity of the polypeptide selected, and the site of injection.

The polypeptides used as an immunogen may be modified as appropriate or administered in an adjuvant in order to increase the peptide antigenicity. Suitable methods increasing antigenicity are well known in the art, and include,
30 for example, coupling the antigen with a heterologous protein (such as globulin or β -galactosidase) or through the inclusion of an adjuvant during immunization.

A preferred method of generating monoclonal antibodies comprises removing spleen cells from the immunized animals, fusing these cells with myeloma cells, such as SP2/0-Ag14 myeloma cells, and allowing them to become monoclonal antibody-producing hybridoma cells. Any one of a number of methods well known in the art may be used to identify the hybridoma cell which produces an antibody with the desired characteristics. These include screening the hybridomas with an ELISA assay, western blot analysis, or radioimmunoassay (Lutz *et al.*, *Exp. Cell Res.* 175:109-124 (1988); Kishimoto *et al.*, *Proc. Natl. Acad. Sci. USA* 87:2244-2248 (1990)). Hybridomas secreting the desired antibodies are cloned and the class and subclass of the secreted antibodies are determined using procedures known in the art (Campbell, A.M., *Monoclonal Antibody Technology: Laboratory Techniques in Biochemistry and Molecular Biology*, Elsevier Science Publishers, Amsterdam, The Netherlands (1984)).

For polyclonal antibodies, antibody-containing antisera is preferably isolated from the immunized animal and is screened for the presence of antibodies with the desired specificity using one of the above-described procedures.

The present invention further provides hybrid cell lines which secrete monoclonal antibodies selective for the peptide agents of Group I and/or Group II. These monoclonal antibodies are capable of wholly or partially neutralizing the activity of the polypeptides or specifically binding to one of the said polypeptides. These monoclonal antibodies can be used for qualitative and/or quantitative measurement or for purification of the polypeptides according to the invention. The present invention therefore also includes test systems which contain the monoclonal antibodies herein described.

Antibodies may be used as an isolated whole antibody, or can be used as a source for generating antibody fragments which contain the antigen binding site of the antibody. Examples of such antibody fragments include, but are not limited to the F_u, the F(ab), the F(ab)₂, fragment, as well as single chain antibodies. Various methods known in the art can be used to generate such fragments without undue experimentation. Recombinant techniques are preferred for generating large quantities of antibodies, antibody fragments and single chain antibodies, as

described, for example, in Pluckthum, *Bio/Technology* 10:163-167 (1992); Carter *et al.*, *Bio/Technology* 10:167-170 (1992); and Mullinax *et al.*, *Biotechniques* 12:864-869 (1992). In addition, recombinant techniques may be used to generate heterobifunctional antibodies.

5 In general, recombinant production of antibodies, antibody fragments or derivatives thereof, uses mRNA encoding an antibody which is isolated from hybridoma cells that produce the desired antibody. This mRNA is then used as a source for generating a cDNA molecule which encodes the antibody, or a fragment thereof. Once obtained, the cDNA may be amplified and expressed
10 according to known methods in a variety of eukaryotic and prokaryotic hosts.

 The present invention further includes derivatives of antibodies (antibody derivatives). As used herein, an "antibody derivatives" contain an antibody of the present invention, or a fragment thereof, as well as an additional moiety which is not normally a part of the antibody. Such moieties may improve the solubility,
15 absorption, biological half-life, etc., of the antibody, decrease the toxicity of the antibody, eliminate or attenuate any undesirable side effect of the antibody, or serve as a detectable marker of the presence of the antibody. Moieties capable of mediating such effects are well known in the art.

 Detectably labeled antibodies constitute a special class of the antibody derivatives of the present invention. An antibody is said to be "detectably labeled"
20 if the antibody, or fragment thereof, is attached to a molecule which is capable of identification, visualization, or localization using known methods. Suitable detectable labels include radioisotopic labels, enzyme labels, non-radioactive isotopic labels, fluorescent labels, toxin labels, affinity labels, chemiluminescent
25 labels and nuclear magnetic resonance contrast agents.

 Illustrative examples of suitable enzyme labels include, but are not limited to, malate dehydrogenase, staphylococcal nuclease, delta-5-steroid isomerase, yeast-alcohol dehydrogenase, alpha-glycerol phosphate dehydrogenase, triose phosphate isomerase, peroxidase, alkaline phosphatase, asparaginase,
30 glucose oxidase, beta-galactosidase, ribonuclease, urease, catalase, glucose-6-phosphate dehydrogenase, glucoamylase, and acetylcholine esterase.

Examples of suitable radioisotopic labels include, but are not limited to, ^3H , ^{111}In , ^{125}I , ^{131}I , ^{32}P , ^{35}S , ^{14}C , ^{51}Cr , ^{57}Co , ^{58}Co , ^{59}Fe , ^{75}Se , ^{152}Eu , ^{90}Y , ^{67}Cu , ^{217}Bi , ^{211}At , ^{212}Pb , ^{47}Sc , ^{109}Pd , etc. ^{111}In is a preferred isotope where *in vivo* imaging is used since it avoids the problem of dehalogenation of the ^{125}I or ^{131}I -labeled monoclonal antibody by the liver. In addition, this radionuclide has a more favorable gamma emission energy for imaging (Perkins *et al.*, *Eur. J. Nucl. Med.* 10:296-301 (1985); Carasquillo *et al.*, *J. Nucl. Med.* 28:281-287 (1987)). For example, ^{111}In coupled to monoclonal antibodies with 1-(p-isothiocyanatobenzyl)-DPTA has shown little uptake in non-tumorous tissues, particularly the liver, and therefore enhances specificity of tumor localization (Esteban *et al.*, *J. Nucl. Med.* 28:861-870 (1987)).

Illustrative examples of suitable non-radioactive isotopic labels include, but are not limited to, ^{157}Gd , ^{55}Mn , ^{162}Dy , ^{52}Tr , and ^{56}Fe .

Illustrative examples of suitable fluorescent labels include, but are not limited to, an ^{152}Eu label, a fluorescein label, an isothiocyanate label, a rhodamine label, a phycoerythrin label, a phycocyanin label, an allophycocyanin label, an o-phthaldehyde label, and a fluorescamine label.

Illustrative examples of suitable toxin labels include diphtheria toxin, ricin, and cholera toxin.

Illustrative examples of chemiluminescent labels include a luminal label, an isoluminal label, an aromatic acridinium ester label, an imidazole label, an acridinium salt label, an oxalate ester label, a luciferin label, a luciferase label, and an aequorin label.

Illustrative examples of nuclear magnetic resonance contrasting agents include paramagnetic heavy metal nuclei such as Gd, Mn, and Fe.

Typical techniques for binding the above-described labels to antibodies are provided by Kennedy *et al.* (*Clin. Chim. Acta* 70:1-31 (1976)), and Schurs *et al.* (*Clin. Chim. Acta* 81:1-40 (1977)), which are both herein incorporated in their entirety by reference. Coupling techniques mentioned in the latter reference include the glutaraldehyde method, the periodate method, the dimaleimide method, and the m-maleimidobenzyl-N-hydroxy-succinimide ester method.

The present invention additionally includes humanized forms of these antibodies. Humanized forms of the antibodies may be produced, for example, by replacing an immunogenic portion of an antibody with a corresponding, non-immunogenic portion (chimeric antibodies) (Better, *et al.*, *Science* 240:1041-1043 (1988)). Alternatively, suitable "humanized" antibodies can be produced by CDR or CEA grafting/substitution as described, for example, in Jones, *et al.*, *Nature* 321:552-525 (1986); Verhoeyan *et al.*, *Science* 239:1534 (1988); and Beidler, *et al.*, *J. Immunol.* 141:4053-4060 (1988).

Another type of the Group II agents of the present invention are peptide agents which are classified as antisense-peptide sequences. Antisense-peptide sequences are short peptides which are specifically designed to bind to a particular amino acid sequence. In general, such antisense peptide agents may be generated using methods known in the art, such as those described, for example, in Hurby *et al.*, "Application of Synthetic Peptides: Antisense Peptides," in *Synthetic Peptides, A User's Guide*, W.H. Freeman, NY, pp. 289-307 (1992) and Kaspczak *et al.*, *Biochemistry* 28:9230-8 (1989).

An additional class of the Group II agents of the invention are natural ligands of APP. As used herein, a natural ligand of APP is defined as any substance which binds to APP, such as soluble fragments of LRP containing the APP binding site. Such soluble fragments may be prepared by any suitable method known to those skilled in the art, such as the method of Davis *et al.*, *Nature* 326:760-765 (1987), which involves deletion of the entire EGF domain. Moreover, soluble forms of the receptor may be formed by inserting a stop codon in front of the region of DNA encoding the cytoplasmic or transmembrane domain (Yokade *et al.*, *J. Cell. Biol.* 117:39 (1992)).

The agents of the present invention may be used *in vitro* and/or *in vivo* to study LRP attachment and to reduce the rate of onset and/or ameliorate the duration and severity of Alzheimer's disease. In addition, the agents of the present invention may be used in qualitative, quantitative and preparative assays and purification procedures to isolate, identify and facilitate the purification of APP.

For *in vivo* use, the agents of the present invention may be provided to a patient as a means of reducing the amount or rate of APP binding to LRP (Hayden, *et al.*, *Antiviral Res.* 9:233-247 (1988)).

5 The present invention therefore provides pharmaceutical compositions comprising a Group I agent and or a Group II agent. These pharmaceutical compositions may be administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, drops or transdermal patch), buccally, or as an oral or nasal spray. As used herein, “pharmaceutically acceptable carrier” is intended to mean a non-toxic solid, 10 semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term “parenteral” as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion. One of ordinary skill will recognize that the choice of a particular mode of administration 15 can be made empirically based upon considerations such as the particular disease state being treated; the type and degree of the response to be achieved; the specific agent or composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration and rate of excretion of the agent or composition; the duration of the treatment; drugs (such as a chemotherapeutic agent) used in combination or coincidental with the specific composition; and like 20 factors well known in the medical arts:

Pharmaceutical compositions of the present invention for parenteral injection may comprise pharmaceutically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions as well as sterile 25 powders for reconstitution into sterile injectable solutions or dispersions just prior to use. Illustrative examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include, but are not limited to, water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), carboxymethylcellulose and suitable mixtures thereof, vegetable oils (such as olive 30 oil), and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials such as lecithin, by the

maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

The compositions of the present invention may also contain adjuvants such as preservatives, wetting agents, emulsifying agents, and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents such as sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of the therapeutic agent or inhibitor, it is desirable to slow the absorption from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use.

Solid dosage forms for oral administration include, but are not limited to, capsules, tablets, pills, powders, and granules. In such solid dosage forms, the

active compounds are preferably mixed with at least one pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents as appropriate.

Solid compositions of a similar type may also be employed as fillers in soft and hard filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Illustrative examples of embedding compositions which can be used include polymeric substances and waxes.

The active agents of Group I and/or Group II can also be in micro-encapsulated form, if appropriate, with one or more of the above-mentioned excipients.

Liquid dosage forms for oral administration include, but are not limited to, pharmaceutically acceptable emulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other

solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethyl formamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

Besides inert diluents, the oral compositions may also contain adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, and tragacanth, and mixtures thereof.

The agent or inhibitor can also be administered in the form of liposomes.

As is known to those skilled in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to the agent or inhibitor, stabilizers, preservatives, excipients, and the like. Preferred lipids are phospholipids and phosphatidyl choline (lecithins), both natural and synthetic. Methods to form liposomes are known in the art. *See, e.g.,* Prescott, ed., *METHODS IN CELL BIOLOGY*, Volume XIV, Academic Press, New York, N.Y. (1976), p. 33 *et seq.*

The agents of the present invention can be formulated according to known methods to prepare pharmaceutically acceptable compositions, whereby these materials, or their functional derivatives, are combined in a mixture with a pharmaceutically acceptable carrier vehicle. Suitable vehicles and their formulation, inclusive of other human proteins, *e.g.,* human serum albumin, are well known in the art. In order to form a pharmaceutically acceptable composition

suitable for effective administration, such compositions will contain an effective amount of one or more agents of the present invention.

Additional pharmaceutical methods may be employed to control the duration of action. Control release preparations may be achieved through the use of polymers to complex or absorb the therapeutic agents of the invention. The controlled delivery may be exercised by selecting appropriate macromolecules (such as polyesters, polyamino acids, polyvinyl, pyrrolidone, ethylenevinylacetate, methylcellulose, carboxymethylcellulose, or protamine sulfate) and methods of incorporation in order to control release. Another possible method to control the duration of action by controlled release preparations is to incorporate antibodies into particles of a polymeric material such as polyesters, polyamino acids, hydrogels, poly(lactic acid) or ethylene vinyl acetate copolymers. Alternatively, instead of incorporating these agents into polymeric particles, it is possible to entrap these materials in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatine-microcapsules and poly(methylmethacrylate) microcapsules, respectively, or in colloidal drug delivery systems, for example, liposomes, albumin microspheres, microemulsions, nanoparticles, and nanocapsules or in macroemulsions.

The pharmaceutical formulations of the present invention are prepared, for example, by admixing the active agent with solvents and/or carriers, optionally using emulsifiers and/or dispersants, whilst if water is used as the diluent, organic solvents may be used as solubilizing agents or auxiliary solvents. As described above, the excipients used include, for example, water, pharmaceutically acceptable organic solvents such as paraffins, vegetable oils, mono- or polyfunctional alcohols, carriers such as natural mineral powders, synthetic mineral powders, sugars, emulsifiers and lubricants.

One of ordinary skill will appreciate that effective amounts of the inventive therapeutic agents can be determined empirically and may be employed in pure form or, where such forms exist, in pharmaceutically acceptable salt, ester or prodrug form. The agonist or antagonist may be administered in compositions in combination with one or more pharmaceutically acceptable excipients. It will be

understood that, when administered to a human patient, the total daily usage of the agents and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgement. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors including the type and degree of the response to be achieved; the specific agent or composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the agent or composition; the duration of the treatment; drugs (such as a chemotherapeutic agent) used in combination or coincidental with the specific composition; and like factors well known in the medical arts.

Techniques of dosage determination are well known in the art for antibody and peptide agents. In general, it is desirable to provide a patient with a dosage of antibody or peptide agent in the range of from about 1 pg/kg to 10 mg/kg (body weight of patient). The therapeutically effective dose can be lowered if the agent of the present invention is additionally administered with another compound. As used herein, one compound is said to be additionally administered with a second compound when the administration of the two compounds is in such proximity of time that both compounds can be detected at the same time in the patient's serum.

For example, satisfactory results are obtained by oral administration of therapeutic dosages on the order of from 0.05 to 10 mg/kg/day, preferably 0.1 to 7.5 mg/kg/day, more preferably 0.1 to 2 mg/kg/day, administered once or, in divided doses, 2 to 4 times per day. On administration parenterally, for example by i.v. drip or infusion, dosages on the order of from 0.01 to 5 mg/kg/day, preferably 0.05 to 1.0 mg/kg/day and more preferably 0.1 to 1.0 mg/kg/day can be used. Suitable daily dosages for patients are thus on the order of from 2.5 to 500 mg p.o., preferably 5 to 250 mg p.o., more preferably 5 to 100 mg p.o., or on the order of from 0.5 to 250 mg i.v., preferably 2.5 to 125 mg i.v. and more preferably 2.5 to 50 mg i.v.

Dosaging may also be arranged in a patient specific manner to provide a predetermined concentration of an agent in the blood, as determined by the RIA

technique. Thus patient dosaging may be adjusted to achieve regular on-going trough blood levels, as measured by RIA, on the order of from 50 to 1000 ng/ml, preferably 150 to 500 ng/ml.

5 The agents of the present invention are intended to be provided to a patient in an amount sufficient to reduce the amount or rate of binding of human APP to LRP. An amount is said to be sufficient to "reduce the amount or rate of APP binding" if the dosage, route of administration, etc. of the agent is sufficient to reduce the amount or rate of APP attachment to the LRP receptor. Such an effect can be assayed, for example, by examining the onset of Alzheimer's disease
10 symptoms occurring *in vivo*, or by correlating *in vitro* blocking studies with predicted *in vivo* efficacy.

The administration of the agents of the present invention may be for either prophylactic or therapeutic purpose. When provided prophylactically, the agent is provided in advance of any Alzheimer's disease symptoms. The
15 prophylactic administration of the agent serves to prevent or reduce the rate of onset of symptoms. When provided therapeutically, the agent is provided at (or shortly after) the onset of the appearance of symptoms of actual disease. The therapeutic administration of the agent serves to reduce the severity and duration of Alzheimer's disease.

20 The present invention further includes the use of the agents of the present invention in diagnostic applications. The Group I agents of the present invention can be used to detect the presence of LRP in a test sample. The Group II agents of the present invention can be used to detect the presence of APP in a test sample.

25 Conditions for incubating an agent with a test sample vary. Incubation conditions will depend on factors such as the type of agent, format, and detection system employed for the assay, as well as the nature of the test sample used in the assay. For example, condition will vary slightly when a whole antibody, a single chain antibody, a F(ab) fragment, or a peptide agent is used. One skilled in the art will recognize that any one of the commonly available immunological assay
30 formats (such as radioimmunoassays, enzyme-linked immunosorbent assays, diffusion based ouchterlony, or rocket immunofluorescent assays) can readily be

adapted to employ the antibodies of the present invention. Examples of such assays can be found in T. Chard, *An Introduction to Radioimmunoassay and Related Techniques*, Elsevier Science Publishers, Amsterdam, The Netherlands (1986); G.R. Bullock *et al.*, *Techniques in Immunocytochemistry*, Academic Press, Orlando, FL Vol. 1 (1982), Vol. 2 (1983), Vol. 3 (1985); and P. Tijssen, P.,
5 *Practice and Theory of Enzyme Immunoassays: Laboratory Techniques in Biochemistry and Molecular Biology*, Elsevier Science Publishers, Amsterdam, The Netherlands (1985).

In one embodiment of the above-described method, the agent of the
10 present invention is immobilized on a solid support for use in the diagnostic assay. Illustrative examples of such solid supports include, but are not limited to, plastics such as polycarbonate, complex carbohydrates such as agarose and sepharose, and acrylic resins, such as polyacrylamide and latex beads. Techniques for coupling agents such as antibodies, peptides and the like to such solid supports are well
15 known in the art, as described, for example, in D.M. Weir *et al.*, *Handbook of Experimental Immunology*, 4th Ed., Blackwell Scientific Publications, Oxford, England, Chapter 10 (1986) and W.D. Jacoby *et al.*, *Meth. Enzym.* 34, Academic Press, N.Y. (1974).

Additionally, one or more of the agents of the present invention which is
20 used in one of the above-described methods can be detectably labelled prior to use, for example, through the use of radioisotopes, affinity labels (such as biotin, avidin, etc.), enzymatic labels (such as horse radish peroxidase, alkaline phosphatase, etc.) fluorescent labels (such as FITC or rhodamine, etc.), paramagnetic atoms, etc. Procedures for accomplishing such labelling are well-
25 known in the art (L.A. Sternberger *et al.*, *J. Histochem. Cytochem.* 18:315 (1970); E.A. Bayer *et al.*, *Meth. Enzym.* 62:308 (1979); E. Engval *et al.*, *Immunol.* 109:129 (1972); and J.W. Goding, *J. Immunol. Meth.* 13:215 (1976)).

The materials used in the inventive assays are ideally suited for the preparation of a kit. For example, the present invention provides a
30 compartmentalized kit to receive in close confinement, one or more containers which comprises:

- a) a first container comprising an agent capable of binding to the LRP binding site; and
- b) one or more other containers comprising one or more of the following: wash reagents and reagents capable of detecting the presence of bound agents from the first container.

5

As used herein, a compartmentalized kit includes any kit in which reagents are contained in separate containers. Illustrative examples of such containers include, but are not limited to, small glass containers, plastic containers or strips of plastic or paper. Particularly preferred types of containers allow the skilled worker to efficiently transfer reagents from one compartment to another compartment such that the samples and reagents are not cross-contaminated and the agents or solutions of each container can be added in a quantitative fashion from one compartment to another. Such containers include, but are not be limited to, a container which will accept the test sample, a container which contains one or more of the agents of the present invention used in the assay, containers which contain wash reagents (such as phosphate buffered saline, Tris-buffers, etc.), and containers which contain the reagents used to detect the bound agent.

10

15

20

The types of detection reagents which can be used in the above described kits include, but are not limited to, labelled secondary agents, or in the alternative, if the primary agent is labelled, enzymatic or agent binding reagents which are capable of reacting with the labelled agent. One skilled in the art will readily recognize that the agents of the present invention can be readily incorporated into one of the established kit formats which are well known in the art.

25

30

A further aspect of the present invention concerns DNA molecules which encode for the polypeptide and antibody agents of Group I and/or Group II. The starting nucleotide molecules can be obtained by the person skilled in the art using known methods. Moreover, the DNA molecules, where the amino acid sequence is known, may be produced synthetically (*e.g.* according to Edge *et al.*, *Nature* 292:756-762 (1981)) or by methods such as PCR (Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press (1989)). The DNA sequences of the present invention include not only the actual

nucleotide sequence used by the organism from which the receptor protein is derived but also includes all degenerate forms which encode a peptide with the desired sequence.

5 The invention includes DNA sequences which have been modified utilizing methods known in the art, such as those generated by mutation, deletion, transposition or addition. The preferred mutations will introduce stop codons within the RAP sequence so that a truncated protein will be generated.

10 The present invention further includes DNA vectors which contain the DNA sequences described above and below. In particular, these may be vectors in which the DNA molecules described are functionally linked to control sequences which allows expression of the corresponding polypeptides. These are preferably plasmids which can be replicated and/or expressed in prokaryotes such as *E. coli* and/or in eukaryotic systems such as yeasts or mammalian cell lines. These vectors may also be mammalian viral vectors which can be replicated and/or
15 expressed in eukaryotes such as mammalian cell lines and in the human patient, as "host," for integration into the cellular genome of the patient and expression as genetic therapy systems.

The invention also includes host organisms transformed with the above vectors. Expression in prokaryotes and eukaryotes may be carried out using
20 techniques known in the art. The DNA sequences according to the invention may be expressed as fusion polypeptides or as intact, native polypeptides. Fusion proteins may advantageously be produced in large quantities. They are generally more stable than the native polypeptide and are easy to purify. The expression of these fusion proteins can be controlled by normal host DNA sequences.

25 For example, the DNA sequences according to the invention can be cloned and expressed as lacZ fusion genes in *E. coli*. A person skilled in the art has a variety of vector systems available for this purpose, *e.g.* the pUR-vector series (U. R  ther and B. M  ller-Hill, *EMBO J.* 2:1791 (1983)). The bacteriophage promoter λp_R may also be used, in the form of the vectors pEX-1 to
30 -3, for expressing large amounts of Cro- β -galactosidase fusion protein (K.K. Stanley and J.P. Luzio, *EMBO J.* 3:1429 (1984)). Analogously, the tac promoter

which can be induced with IPTG can also be used, for example in the form of the pROK-vector series (CLONTECH Laboratories).

The prerequisite for producing intact native polypeptides using *E. coli* is the use of a strong, regulatable promoter and an effective ribosome binding site. Promoters which may be used for this purpose include the temperature sensitive bacteriophage λ p_L-promoter, the tac-promoter inducible with IPTG or the T7-promoter. Numerous plasmids with suitable promoter structures and efficient ribosome binding sites have been described, such as for example pKC30 (λ p_L; Shimatake and Rosenberg, *Nature* 292:128 (1981), pKK173-3 (tac, Amann and Brosius, *Gene* 40:183 (1985)) or pET-3 (T7-promoter (Studier and Moffat, *J. Mol. Biol.* 189:113 (1986)).

A number of other suitable vector systems for expressing the DNA according to the invention in *E. coli* are known from the prior art and are described, for example, in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press (1989)).

Suitable *E. coli* strains which are specifically tailored to a particular expression vector are known to those skilled in the art (Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press (1989)). The experimental performance of the cloning experiments, the expression of the polypeptides in *E. coli* and the working up and purification of the polypeptides are known and are described, for example, in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press (1989). In addition to prokaryotes, eukaryotic microorganisms such as yeast may also be used.

For expression in yeast, the plasmid YRp7 (Stinchcomb *et al.* *Nature* 282:39 (1979); Kingsman *et al.*, *Gene* 7:141 (1979); Tschumper *et al.*, *Gene* 10:157 (1980)) and the plasmid YEp13 (Bwach *et al.*, *Gene* 8:121-133 (1979)) are used, for example. The plasmid YRp7 contains the TRP1-gene which provides a selection marker for a yeast mutant (*e.g.* ATCC No. 44076) which is incapable of growing in tryptophan-free medium. The presence of the TRP1 defect as a characteristic of the yeast strain used then constitutes an effective aid to detecting

transformation when cultivation is carried out without tryptophan. The same is true with the plasmid YEp13, which contains the yeast gene LEU-2, which can be used to complete a LEU-2-minus mutant.

Other suitable marker genes for yeast include, for example, the URA3- and HIS3-gene. Preferably, yeast hybrid vectors also contain a replication start and a marker gene for a bacterial host, particularly *E. coli*, so that the construction and cloning of the hybrid vectors and their precursors can be carried out in a bacterial host. Other expression control sequences suitable for expression in yeast include, for example, those of PHO3- or PHO5-gene.

Other suitable promoter sequences for yeast vectors contain the 5'-flanking region of the genes of ADH I (Ammerer, *Methods of Enzymology* 101:192-210 (1983)), 3-phosphoglycerate kinase (Hitzeman *et al.*, *J. Biol. Chem.* 255:2073 (1980)) or other glycolytic enzymes (Kawaski and Fraenkel, *BBRC* 108:1107-1112 (1982)) such as enolase, glyceraldehyde-3-phosphate-dehydrogenase, hexokinase, pyruvate-decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, phosphoglucose-isomerase and glucokinase. When constructing suitable expression plasmids, the termination sequences associated with these genes may also be inserted in the expression vector at the 3'-end of the sequence to be expressed, in order to enable polyadenylation and termination of the mRNA.

Other promoters are the promoter regions of the genes for alcohol dehydrogenase-2, isocytochrome C, acid phosphatase and enzymes which are responsible for the metabolism of maltose and galactose. Promoters which are regulated by the yeast mating type locus, such as promoters of the genes BARI, MF α 1, STE2, STE3, STE5 can be inserted in temperature regulated systems by the use of temperature-dependent sir mutations. (Rhine, Ph.D. Thesis, University of Oregon, Eugene, Oregon (1979); Herskowitz and Oshima, *The Molecular Biology of the Yeast Saccharomyces*, Cold Spring Harbor Laboratory Press, part I, pp. 181-209 (1981)). Generally, however, any vector which contains a yeast-compatible promoter and origin replication and termination sequences is suitable. Thus, hybrid vectors which contain sequences homologous to the yeast 2 μ

plasmid DNA may also be used. Such hybrid vectors are incorporated by recombination within the cells of existing 2 μ -plasmids or replicate autonomously.

In addition to yeasts, other eukaryotic systems may, of course, be used to express the polypeptides according to the invention. Since post-translational modifications such as disulphide bridge formation, glycosylation, phosphorylation and/or oligomerization are frequently necessary for the expression of biologically active eukaryotic proteins by means of recombinant DNA, it may be desirable to express the DNA according to the invention not only in mammalian cell lines but also insect cell lines.

Functional prerequisites of the corresponding vector systems comprise, in particular, suitable promoter, termination and polyadenylation signals as well as elements which make it possible to carry out replication and selection in mammalian cell lines. For expression of the DNA molecules according to the invention it is particularly desirable to use vectors which are replicable both in mammalian cells and also in prokaryotes such as *E. coli*.

Vectors derived from viral systems such as SV40, Epstein-Barr-virus, etc., include, for example, pTK2, pSV2-dhfv, pRSV-neo, pKO-neo, pHyg, p205, pHEBo, etc. (Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press, N.Y. (1989)).

After transformation in suitable host cells, *e.g.* CHO-cells, corresponding transformed cells may be obtained with the aid of selectable markers (thymidine-kinase, dihydrofolate-reductase etc.) and the corresponding polypeptides are isolated after expression. The host cells suitable for the vectors are known, as are the techniques for transformation (micro-injection, electroporation, calcium phosphate method, etc.) as described, for example, in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press, N.Y. (1989).

For cloning corresponding DNA fragments in prokaryotic or eukaryotic systems, the selected vector may cut, for example, with a restriction endonuclease and, optionally after modification of the linearized vector thus formed, an expression control sequence equipped with corresponding restriction ends is inserted. At the 3'-end (in the direction of translation) the expression control

sequence contains the recognition sequence of a restriction endonuclease, so that the vector already containing the expression control sequence is digested with the said restriction enzyme and the DNA molecule according to the invention, provided with ends which fit, can be inserted. It is advantageous to cleave the vector which already contains the expression control sequence with a second restriction endonuclease inside the vector DNA and to insert the DNA molecule provided with the correct ends into the vector fragment produced. The techniques required are described, for example, by Sambrook *et al.* *Molecular Cloning: A Laboratory Manual* Cold Spring Harbor Press. N.Y. (1989).

Apart from the DNA molecules specified, the invention also relates to processes for preparing the vectors described herein, particularly expression vectors. These vectors are characterized in that a DNA provided with corresponding ends and coding for a functional derivative or a fragment of the LRP receptor is inserted into a vector DNA cut with restriction endonucleases and containing the expression control sequences described by way of example, in such a way that the expression control sequences regulate the expression of the DNA inserted. The peptides and antibody agents of the present invention which are obtained by the expression of recombinant DNA or from the native receptor molecule may, of course, also be derivatized by chemical or enzymatic processes.

The present invention is described in further detail in the following non-limiting examples.

Example 1: ¹²⁵I-labeled APP₇₇₀ Degradation by LRP-expressing Fibroblasts

Wells containing 2×10^5 LRP-expressing (MEF) or LRP-deficient (PEA13) cells were incubated with various concentrations (0.37 - 10 nM) of ¹²⁵I-APPs770 (Panels A and B) or ¹²⁵I-APPs695 (Panels C and D) for 10 h at 37 °C in the absence of competitor (●) or in the presence of either 1 μM RAP (○), or 100 μg/ml heparin (□). The cellular degradation of ¹²⁵I-labeled APP₇₇₀ by LRP-expressing MEF cells is shown in panel A and by LRP-deficient PEA13 cells in panel B. The cellular degradation of ¹²⁵I-APP₆₉₅ by LRP-expressing cells is depicted in panel C and by LRP-deficient cells in panel D. Plotted values

represent means of duplicate determinations. *Inset* to panel *B*, autoradiogram of ^{125}I -labeled APP₇₇₀ (lane 1) and APP₆₉₅ (lane 2). The migration position of molecular mass standards are indicated on the right in kDa.

5 ***Example 2: LRP-deficient cells internalize and degrade ^{125}I -LDL, but not ^{125}I - $\alpha_2\text{M}$***

Wells containing 2×10^5 LRP-expressing (●) or LRP-deficient (○) mouse fibroblasts were incubated with ^{125}I -LDL (8 $\mu\text{g/ml}$) or ^{125}I - $\alpha_2\text{M}^*$ (1.4 nM) for selected time intervals. Specific internalization and degradation was
10 determined by incubation of ^{125}I -ligand in the presence of a 200-fold excess unlabeled ligand. Shown are the amounts of ^{125}I -LDL internalized (*panel A*) and degraded (*panel B*), ^{125}I - $\alpha_2\text{M}^*$ internalized (*panel C*) and degraded (*panel D*). The plotted values represent means of duplicate determinations.

15 ***Example 3: Time course of the cellular internalization and degradation of ^{125}I -APP₇₇₀***

Mouse fibroblasts (2×10^5 cells per well) were incubated for selected times with ^{125}I -APP₇₇₀ (1 nM) in the absence of competitor (●) or in the presence of either 1 μM RAP (○), or 100 $\mu\text{g/ml}$ heparin (□). The amount of
20 radioactivity internalized and degraded was determined at the indicated time intervals. Shown is the amount of internalization and degradation of ^{125}I -APP₇₇₀ by LRP-expressing MEF cells (*panels A and B, respectively*) and the internalization and degradation of ^{125}I -APP₇₇₀ by LRP-deficient PEA13 cells (*panels C and D, respectively*). The plotted values are means of duplicate
25 determinations.

Example 4: LRP-antibodies inhibit the degradation of ^{125}I -APP₇₇₀ by mouse fibroblasts

Mouse fibroblasts (2×10^5 cells/well) were incubated for 18 h at 37 °C
30 with ^{125}I -APP₇₇₀ (10 nM) in the absence of competitor or in the presence of either heparin (100 $\mu\text{g/ml}$), chloroquine (0.1 mM), RAP (1 μM), affinity purified

anti-LRP IgG (100 µg/ml), or control IgG against the cytoplasmic domain of LRP (100 µg/ml). Panels show the amount of ¹²⁵I-APP₇₇₀ degraded by mouse fibroblasts expressing LRP (A) or ones deficient in LRP expression (B). The data shown are means of duplicate determinations ± S. E.

5

Example 5: LRP binds directly to APP₇₇₀ in ELISA

In panel A, zinc-coated wells (○) or zinc-coated wells incubated with APP₇₇₀ (●) were incubated with the APP₇₇₀ monoclonal antibody 7H5. In panel B, increasing concentrations of LRP (0.14-300 nM) were incubated for 18 h at 4°C with zinc-coated wells (○) or zinc-coated wells with captured APP₇₇₀ (●). Panel C, LRP (25 nM) was incubated with zinc-coated wells (○) or zinc-coated wells with captured APP₇₇₀ (●) in the presence of increasing concentrations of RAP (0.2 - 450 nM). Bound receptor was detected with the monoclonal LRP antibody 8G1. In the absence of LRP there was no binding of 8G1 IgG to APP₇₇₀-coated wells. Plotted values represent means of duplicate determinations.

15

Example 6: ¹²⁵I-APP₇₇₀:Factor IXa complex is degraded by LRP-expressing cells but not by LRP-deficient cells

Wells containing 2 x 10⁵ LRP-expressing (MEF) or LRP-deficient (PEA13) cells were incubated with ¹²⁵I-APP₇₇₀ (5 nM) or ¹²⁵I-APP₇₇₀:FIXa (3 nM) for 18 h at 37°C in the absence of competitor or in the presence of 1 µM RAP. The amount of ¹²⁵I-labeled APP₇₇₀ or ¹²⁵I-APP₇₇₀:Factor IXa degraded by LRP-expressing MEF cells is shown in panel A and that by LRP-deficient PEA13 cells in panel B. Plotted values represent means of duplicate determinations.

20

25

Example 7: Effect of apoE isoforms on the degradation of ¹²⁵I-APP₇₇₀ by mouse fibroblasts

Mouse fibroblasts (2 x 10⁵) were incubated for 18 h at 37°C with ¹²⁵I-APP/PN-2 (8.8 nM) in the absence of competitor or in the presence of either RAP (1 µM), 10 µg/ml apoE3/βVLDL, 10 µg/ml apoE4/βVLDL, or 10 µg/ml βVLDL.

30

The concentration of apoE-enriched β VLDL represent the β VLDL protein before enrichment with apoE. Panels show the amount of ^{125}I -APP₇₇₀ degradation by mouse fibroblasts expressing LRP (A) or deficient in LRP expression (B). The data shown are means of duplicate determinations \pm S. E. The inhibition was statistically significant for both apoE3- β VLDL and apoE4- β VLDL, and for RAP F(4,15)=21.65, $p=.0001$; Fisher protected least significant difference post hoc test showed that both apoE3- β VLDL and apoE4- β VLDL were different from β VLDL alone ($p<0.05$ for both) but not from each other. RAP was also significantly different than all other categories ($p<0.001$).

It will be apparent to those skilled in the art that various modifications and variations can be made in the methods of the present invention without departing from the spirit and scope of the invention. Thus, it is intended that the present invention cover the modifications and variations of this invention provided they come within the scope of the appended claims and their equivalents.

The disclosures of all patents, patent applications, and publications referred to above are hereby expressly incorporated herein by reference.

What is claimed is:

1. A method for preventing catabolism of amyloid β -precursor protein in a mammalian cell which comprises contacting said cell with an agent that reduces the amount or rate of binding of amyloid β -precursor protein (APP) with one or more members of the low density lipoprotein (LDL) receptor family.
5
2. The method according to claim 1, wherein said member of the low density lipoprotein receptor family is the low density lipoprotein receptor-related protein (LRP).
10
3. The method according to claim 2, wherein said agent is an agent that which binds to the amyloid β -precursor protein binding site on the low density lipoprotein receptor-related protein.
15
4. The method according to claim 2, wherein said agent is an agent that binds to the low density lipoprotein receptor-related protein binding site found on amyloid β -precursor protein.
- 20 5. The method according to claim 3, wherein said agent is a functional derivative of amyloid β -precursor protein (APP) or receptor associated protein (RAP).
- 25 6. The method according to claim 4, wherein said agent is an antibody or an antibody fragment containing the antigen binding domain that binds to the LRP binding site found on APP.
7. The method according to claim 4, wherein said agent is a functional derivative of LRP.
30

8. A test kit for determining whether an agent reduces the amount or rate of binding of amyloid β -precursor protein (APP) with the low density lipoprotein receptor-related protein (LRP), comprising a carrier means having in close confinement therein one or more container means at least one of which contains an antibody that binds to the LRP binding site found on APP.
9. A process for preparing antibodies that bind to the low density lipoprotein receptor-related protein (LRP) binding site found on amyloid β -precursor protein (APP), which comprises
- immunizing a host animal with an agent that binds to the LRP binding site on APP;
 - fusing the B-lymphocytes of these host animals with myeloma cells; and
 - subcloning and cultivating a hybrid cell line secreting a monoclonal antibody that binds to the LRP binding site found on APP.
10. A method for reducing the rate of onset or the severity of Alzheimer's disease, which comprises administering to an animal an agent which binds to the low density lipoprotein receptor-related protein (LRP) binding site on the amyloid β -precursor protein (APP) and/or which binds to the APP binding site on the LRP, in an amount effective to reduce the rate of APP attachment to the LRP.
11. A pharmaceutical composition for therapeutic treatment of Alzheimer's disease, comprising at least one agent selected from the group consisting of an agent which binds to the low density lipoprotein receptor-related protein (LRP) binding site on the amyloid β -precursor protein (APP), an agent which binds to the APP binding site on the LRP, and the native receptor molecule of the LDL-receptor family; and

a pharmaceutically acceptable carrier.

- 5
12. A method for identifying substances which inhibit the binding of a ligand (RAP) or amyloid β -precursor protein (APP) to a protein of the low density lipoprotein receptor family, comprising the steps of:
- a) incubating the receptor, or a soluble form of the receptor, with RAP or APP in the presence of a potential inhibitor substance; and
- 10 b) determining the extent of binding of RAP or APP to the receptor or receptor fragment.
13. A method for detecting receptors of the low density lipoprotein receptor family, comprising the steps of:
- 15 a) incubating a substance derived from a fragment of receptor associated protein (RAP) or amyloid β -precursor protein (APP) which contains a binding activity for the receptor with a sample; and
- b) determining the extent of binding of the RAP or APP material to the sample.
- 20
14. A method for supplying a therapeutically active substance into a cell, comprising the steps of:
- 25 a) coupling a therapeutically active substance with a fragment of receptor associated protein (RAP) or amyloid β -precursor protein (APP) with a binding activity on the low density lipoprotein receptor; and
- b) contacting said therapeutically active substance coupled to said fragment of RAP or APP with said cell under conditions sufficient to bind said fragment of RAP or APP to said receptor in a manner effective to introduce said therapeutically active substance into said cell.
- 30

1 / 11




FIG. 1A





FIG. 1B



2/11




FIG. 1D




FIG. 1C




FIG. 2B




FIG. 2A

fmoles $^{125}\text{I-}\alpha_2\text{M}^*$ degraded/ 10^5 cells




FIG.2D




FIG.2C

5/11




FIG. 3B




FIG. 3A

6/11




FIG. 3D




FIG. 3C

7 / 11




FIG. 4B




FIG. 4A

8/11




FIG. 5B

FIG. 5A

9/11




FIG. 5C

11/11




FIG. 6B




FIG. 6A

11/11




FIG. 7B




FIG. 7A

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/12686

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/130.1, 141.1, 143.1; 435/7.21, 70.21; 436/547, 548; 514/2; 530/345, 350, 402, 387.1, 388.1, 388.22

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched


Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Cell, Volume 6, issued 1995, Kounnas et al, "LDL receptor-related protein (LRP) binds secreted β -amyloid precursor protein and mediates its degradation", page 327A, see abstract 1901.	1-7 and 10-11
Y, P	Fibrinolysis, Volume 10, issued June 1996, Horn et al, "Recombinant fragments of the second cluster of complement-type repeats of LRP mediate high affinity binding of different ligands as determined by surface plasmon resonance", abstract 61, see entire abstract.	1-7 and 10-11

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

•	Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A"	document defining the general state of the art which is not considered to be of particular relevance		
"E"	earlier document published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means		
"P"	document published prior to the international filing date but later than the priority date claimed	"Z"	document member of the same patent family

Date of the actual completion of the international search 15 OCTOBER 1996	Date of mailing of the international search report 08 NOV 1996
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer PATRICIA A. DUFFY  Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

 International application No.
 PCT/US96/12686

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Neuroscience, Volume 65, Number 4, issued 1995, Lorent et al, "Expression in mouse embryos and in adult mouse brain of three members of the amyloid precursor protein family, of the alpha-2-macroglobulin receptor/low density lipoprotein receptor-related protein and of its ligands apolipoprotein E, lipoprotein lipase, alpha-2-macroglobulin and the 40,000 molecular weight receptor-associated protein", pages 1009-1025, see entire document.	1-7 and 10-11
A	Journal of Lipid Research, Volume 36, Number 9, issued 1995, Raffai et al, "Molecular characterization of two monoclonal-antibodies specific for the LDL receptor-binding site of human apolipoprotein E", pages 1905-1918, see entire document.	8 and 9
Y	Annals of Neurology, Volume 37, Number 2, issued February 1995, Rebeck et al, "Multiple, diverse senile plaque-associated proteins are ligands of an apolipoprotein E receptor, the α_2 -macroglobulin receptor/low-density-lipoprotein receptor-related protein", pages 211-217, see entire document.	1-7 and 10-11
X,P — Y,P	Cell, Volume 82, No. 2, issued 28 July 1995, Kounnas et al, "LDL Receptor-Related Protein, a Multifunctional ApoE Receptor, Binds Secreted β -Amyloid Precursor Protein and Mediates Its Degradation", pages 331-340, see entire document.	1-9 ——— 10-11

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/12686

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-11

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/12686

A. CLASSIFICATION OF SUBJECT MATTER:
IPC (6):

A61K 37/18, 38/00, 39/395; C07K 1/00, 14/00, 16/00, 17/00; C12P 21/04; G01N 33/53, 33/567

A. CLASSIFICATION OF SUBJECT MATTER:
US CL :

424/130.1, 141.1, 143.1; 435/7.21, 70.21; 436/547, 548; 514/2; 530/345, 350, 402, 387.1, 388.1, 388.22

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, DIALOG, MEDLINE, BIOSYS, CAB ABSTRACTS, EMBASE, JAPMO, DERWENT WPI, DERWENT BIOTECHNOLOGY ABSTRACTS.

search terms: low density related lipoprotein receptor, LRP, receptor associated protein, RAP, amyloid, amyloid precursor protein, APP, antibody, immunoglobulin, monoclonal, screen, detect, conjugate, analog, agonist, antagonist.

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

Group I, claim(s) 1-11, drawn to methods for preventing the catabolism of amyloid β -precursor protein in vivo and in vitro by contacting a cell with an agent which reduces the amount or rate of binding of the amyloid β -precursor protein with one or more members of the low density lipoprotein receptor family.

Group II, claim(s) 12, drawn to a method for identifying substances which inhibit the binding of a ligand (RAP) or APP to a protein of the low density lipoprotein receptor family.

Group III, claim(s) 13, drawn to a method for detecting receptors of the low density lipoprotein receptor family.

Group IV, claim(s) 14, drawn to a method for supplying a therapeutically active substance into a cell.

The special technical features are as follows:

Group I - agents which reduce the rate of binding of the amyloid β -precursor protein with one or more members of the low density lipoprotein receptor family.

Group II - method of screening for agents which inhibit the bind of RAP or APP to one or more members of the low density lipoprotein receptor family, this is a second method which relies on the inhibition of binding of APP to the one or more members of the low density lipoprotein receptor family.

Group III - method for detecting receptors of the low density lipoprotein receptor family using RAP or APP as specific binding agents, and thus RAP and APP are the special technical features of this method.

Group IV - an agent comprised of a therapeutically active substance coupled to a fragment of RAP or APP, thus the coupled agent is the special technical feature. The inventions listed as Groups I, II, III and IV do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: the groups fail to share the same or similar special technical feature as defined directly above. Because the groups fail to share a special technical feature, the claims are not so linked by a special technical feature within the meaning of PCT Rule 13.2 so as to form a single inventive concept.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☒ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.